

THE SINGULAR COHOMOLOGY RING OF A UNIFORM MATROID: COMBINATORICS AND LEFSCHETZ PROPERTIES

KYLE BINDER

ABSTRACT. The singular cohomology ring of a matroid is an algebraic invariant which generalizes the Chow ring of a matroid. We study combinatorial and Lefschetz properties of the singular cohomology ring of a uniform matroid. Combinatorially, we construct an explicit basis for the singular cohomology ring in terms of Koszul homology. From this basis we derive multiple formulas for the Hodge numbers of the cohomology ring that recover and extend known formulas for the Chow polynomial of a uniform matroid. We also use this basis to show that the singular cohomology ring of a uniform matroid satisfies the *quasi-projective strong Lefschetz property*—a slight weakening of the Hard Lefschetz property found in the Chow ring of a matroid.

1. INTRODUCTION

The *singular cohomology ring* of a loopless matroid M was introduced in [Bin26] as an extension of the well-studied *Chow ring* of a matroid. Both algebraic invariants are defined via the smooth toric variety X_{Σ_M} associated to the *Bergman fan* Σ_M of the matroid. The Chow ring of M is the Chow ring $A^\bullet(X_{\Sigma_M})$ of the toric variety, while the singular cohomology ring of M is the singular cohomology ring $H^\bullet(X_{\Sigma_M})$ of the toric variety. (All Chow and singular cohomology rings here are taken with rational coefficients.) As the cycle class map $A^\bullet(X_{\Sigma_M}) \hookrightarrow H^{2\bullet}(X_{\Sigma_M})$ is injective for smooth toric varieties [Tot14], the singular cohomology ring of M contains the Chow ring as a subring.

At this stage there is still much to be discovered about these singular cohomology rings. In general, we have vanishing results for the cohomology and a computation of the top-weight cohomology [Bin26, Theorem 6.6], but we do not have formulas for the Betti numbers or even a description for generators of the singular cohomology ring. The algebraic structure of the singular cohomology ring is also not understood. One of the salient properties of the Chow ring of a matroid is that it satisfies the *Kähler package*, and in particular, has the Hard Lefschetz property for multiplication by a generic linear form. It is not known whether the singular cohomology ring of a matroid also has such properties.

The goal of this paper is to address these questions in the context of uniform matroids. Here, we construct an explicit, combinatorial basis for the singular cohomology ring of a uniform matroid in terms of Koszul homology, use these bases to produce explicit formulas for the Betti numbers (and the finer *Hodge numbers*), and show that, even though Hard Lefschetz fails, the singular cohomology ring of a uniform matroid satisfies the *quasi-projective strong Lefschetz property*.

Our results also provide useful information about the Chow rings of uniform matroids. For example, the basis we construct for the singular cohomology ring of a uniform matroid restricts to a new and combinatorially rich basis for the Chow ring. By studying the combinatorics of the basis elements, we produce combinatorial proofs of several formulas for the Chow polynomials of uniform matroids due to Hoster [Hos26], Hameister, Rao, and Simpson [HRS21], and Ferroni, Matthews, Stevens, and Vecchi [Fer+24]. This suggests that the singular cohomology ring may be a useful tool in studying the Chow ring of a general matroid.

Let us now describe our main results in more detail.

1.1. Combinatorics of the singular cohomology ring of a uniform matroid. The Chow ring of a loopless matroid on the ground set $[n] = \{1, \dots, n\}$ has a well-known combinatorial presentation given by Feichtner and Yuzvinsky [FY04]:

$$(1) \quad A^\bullet(X_{\Sigma_M}) = \frac{\mathbb{Q}[\Sigma_M]}{\left(\sum_{i \in F} x_F - \sum_{j \in G} x_G : i, j \in [n]\right)}.$$

Here, $\mathbb{Q}[\Sigma_M]$ is the *Stanley–Reisner ring* of the Bergman fan. This is the ring generated by indeterminates x_F indexed by the proper, non-empty flats of M and subject only to the relation $x_F x_G = 0$ whenever F and G are incomparable. Feichtner and Yuzvinsky also produce a basis for the Chow ring using Gröbner bases.¹

The singular cohomology ring similarly has a combinatorial description. Let $N_{\mathbb{Q}}^{\vee}$ be the character lattice of X_{Σ_M} tensored with \mathbb{Q} . This vector space is spanned by elements $\ell_i - \ell_j$ for $i, j \in [n]$. We then have the bigraded *Koszul complex*

$$K_i(\mathbb{Q}[\Sigma_M])_j = \mathbb{Q}[\Sigma_M]_{j-i} \otimes \bigwedge^i N_{\mathbb{Q}}^{\vee}$$

whose differential satisfies the Leibniz rule, with $d(x_F) = 0$ and $d(\ell_i - \ell_j) = \sum_{i \in F} x_F - \sum_{j \in G} x_G$. Taking homology produces an isomorphism of rings

$$H_\bullet(K(\mathbb{Q}[\Sigma_M])) \cong H^\bullet(X_{\Sigma_M}).$$

Moreover, the bigrading of the Koszul complex induces the weight filtration on $H^\bullet(X_{\Sigma_M})$ [Web04]

$$H_i(K(\mathbb{Q}[\Sigma_M]))_j \cong \text{Gr}_{2j}^W H^{2j-i}(X_{\Sigma_M}),$$

and the Chow ring corresponds to $H_0(K(\mathbb{Q}[\Sigma_M]))$ and thus the associated graded $\text{Gr}_{2j}^W H^{2j}(X_{\Sigma_M})$.

Our first result in this paper is an explicit basis for the singular cohomology of uniform matroids. We describe this basis using what we call *retral* and *weakly retral* chains of flats.

Theorem 1.1. *The set*

$$\left\{ \prod_{F \in \mathcal{F}} x_F : \mathcal{F} \text{ is retral} \right\} \cup \left\{ \prod_{F \in \mathcal{F}} x_F \otimes \xi : \mathcal{F} \text{ is weakly retral and } \xi \text{ is admissible to } \mathcal{F} \right\}$$

is a basis of $H_\bullet(K(\mathbb{Q}[\Sigma_{U_{r,n}}])) \cong H^\bullet(X_{\Sigma_{U_{r,n}}})$ as a \mathbb{Q} -vector space.

We construct this basis in Section 3 using Gröbner theory and an induction on rank. One of the novelties of this result is that for $r < n$ the toric variety $X_{\Sigma_{U_{r,n}}}$ is not proper. To this point, most work on the cohomology of toric varieties has focused on the compact (and potentially singular) case.

Restricting to $H_0(K(\mathbb{Q}[\Sigma_{U_{r,n}}]))$ produces a new basis for the Chow ring of a uniform matroid.

Corollary 1.2. *The set*

$$\left\{ \prod_{F \in \mathcal{F}} x_F : \mathcal{F} \text{ is retral} \right\}$$

is a basis of $A^\bullet(X_{\Sigma_M})$ as a \mathbb{Q} -vector space.

¹In their paper, Feichtner and Yuzvinsky use a slightly different presentation than what we use here. They include an indeterminate for the maximal flat and quotient by the linear forms $\sum_{i \in F} x_F$.

Our second main result is to produce combinatorial formulas for the *Hodge numbers*—the dimensions of the associated graded of the weight filtration—of the singular cohomology of $U_{r,n}$. This is the subject of Section 4.

We will package the Hodge numbers of the singular cohomology of a matroid M in two ways. The first is in terms of a Macaulay2 Betti diagram²

	0	1	2	3	...
0	$\beta_{0,0}$	$\beta_{1,1}$	$\beta_{2,2}$	$\beta_{3,3}$...
1	$\beta_{0,1}$	$\beta_{1,2}$	$\beta_{2,3}$	$\beta_{3,4}$...
2	$\beta_{0,2}$	$\beta_{1,3}$	$\beta_{2,4}$	$\beta_{3,5}$...
⋮	⋮	⋮	⋮	⋮	⋱

where $\beta_{i,j} = \dim H_i(K(\mathbb{Q}[\Sigma_M]))_j = \dim \text{Gr}_{2j}^W H^{2j-i}(X_{\Sigma_M})$. The first column of the Betti diagram displays the Betti numbers of the Chow ring.

The second packaging of Hodge numbers is with the bivariate *Hodge–Poincaré polynomial*

$$\text{HPoin}_M(w, x) = \sum_{i,j} \dim H_i(K(\mathbb{Q}[\Sigma_M]))_j \cdot w^i x^j = \sum_{i,j} \dim \text{Gr}_{2j}^W H^{2j-i}(X_{\Sigma_M}) \cdot w^i x^j.$$

It will often be convenient to break the Hodge–Poincaré polynomial into univariate *refined Hodge–Poincaré polynomials*

$$\underline{H}_M^i(x) = \sum_j \dim H_i(K(\mathbb{Q}[\Sigma_M]))_j \cdot x^{j-i},$$

whence

$$\text{HPoin}_M(w, x) = \sum_{i=0}^{n-r} \underline{H}_M^i(x) \cdot w^i x^i.$$

Note that $\underline{H}_M^i(x)$ is the generating function for the i th column of the Betti diagram for the singular cohomology ring and that $\underline{H}_M^0(x)$ is the *Chow polynomial* (or *Hilbert–Poincaré series* of the Chow ring) of M , which we will also denote by $\underline{H}_M(x)$.

The Chow polynomials of uniform matroids are well-studied and have numerous combinatorial formulas. The following is only a partial list of results. (See Section 4 for any undefined notation.)

Theorem 1.3. *Let $U_{r,n}$ be the uniform matroid of rank $r \geq 1$ on ground set $[n]$. Then*

(1) [Hos26, Theorem 3]

$$\underline{H}_{U_{r,n}}(x) = \sum_{\substack{R \subseteq \{0, \dots, r-1\} \\ 0 \in R}} \binom{n}{\Delta R} \cdot x^{|\Delta R|-1}.$$

(2) [HRS21, Theorem 5.1]

$$x \cdot \underline{H}_{U_{n-1,n}}(x) = d_n(x).$$

(3) [Fer+24, Theorem 1.9]

$$\underline{H}_{U_{r,n}}(x) = \sum_{j=0}^{r-1} \binom{n}{j} d_j(x) \cdot (1 + x + \dots + x^{r-j-1}).$$

²This is precisely the Macaulay2 Betti diagram for $\mathbb{Q}[\Sigma_M]$ as an $\text{Sym}(N_{\mathbb{Q}}^{\vee})$ -algebra.

(4) [Fer+24, Equation (13)]

$$\underline{H}_{U_{r,n}}(x) = \sum_{j=0}^{r-1} \sum_{m=0}^{r-j-1} (-1)^m \binom{n}{j} \binom{n-j-1}{m} A_j(x) \cdot x^{r-m-j-1}.$$

From the combinatorics of retral and weakly retral chains, we give combinatorial proofs for (1)–(3) and rederive (4).

We also produce analogous formulas for the other refined Hodge–Poincaré polynomials.

Theorem 1.4. *Let $U_{r,n}$ be the uniform matroid of rank $r \geq 1$ on ground set $[n]$. For $i \geq 1$,*

(1) [Theorem 4.9]

$$\underline{H}_{U_{r,n}}^i(x) = \sum_{\substack{R \subseteq \{0, \dots, r-1\} \\ 0, r-1 \in R}} \sum_{\ell=1}^{n-r-i+1} \binom{n}{\Delta R} \binom{n - \lfloor R \rfloor - \ell}{r - \lfloor R \rfloor - 1} \binom{n-r-\ell}{i-1} \cdot x^{|R|-1}.$$

(2) [Theorem 4.15]

$$\underline{H}_{U_{r,n}}^i(x) = \sum_{j=0}^{r-1} \sum_{\ell=1}^{n-r-i+1} \binom{n}{j} \binom{n-j-\ell}{r-j-1} \binom{n-r-\ell}{i-1} d_j(x) \cdot x^{r-j-1}.$$

(3) [Proposition 4.18]

$$\underline{H}_{U_{r,n}}^i(x) = (-1)^{r-1} \binom{n-1}{i+r-1} \cdot x^{r-1} + \sum_{j=1}^{r-1} (-1)^{r+1-j} \binom{n}{j} \binom{n-j-1}{i+r-j-1} A_j(x) \cdot x^{r-j}.$$

1.2. Lefschetz properties in the singular cohomology ring of a uniform matroid. One of the miraculous properties of the Chow ring of a matroid is that it acts like the cohomology ring of a smooth projective variety even though the variety X_{Σ_M} is often non-proper and thus only quasi-projective. Specifically, Adiprasito, Huh, and Katz [AHK18] proved that $A^\bullet(X_{\Sigma_M})$ satisfies the *Kähler package* of Poincaré duality, Hard Lefschetz, and the Hodge–Riemann relations. A natural question is the following.

Question 1.5. *Does the singular cohomology ring of a matroid satisfy anything like the Kähler package?*

Unfortunately, the Kähler package itself quickly fails for the singular cohomology rings of matroids. By [Bin26, Theorem 6.6], the Betti diagram for the singular cohomology of a simple rank r matroid M on ground set $[n]$ takes the form

	0	1	⋯	$n-r-1$	$n-r$
0	1	0	⋯	0	0
1	*	*	⋯	*	*
⋮	⋮	⋮	⋱	⋮	⋮
$r-2$	*	*	⋯	*	*
$r-1$	1	*	⋯	*	$\mu(M)$

where asterisks indicate the locations of potentially non-zero entries. The dimension of the top-degree singular cohomology is $\mu(M)$, the *Möbius invariant* of M . For loopless matroids, $\mu(M) \geq 1$, and often this inequality is strict. This immediately implies that Poincaré duality fails in general, as the dimension the top-degree cohomology is not 1. Hard Lefschetz, which implies Poincaré duality,

thereby fails as well. Finally, as there is no degree map on the top-degree cohomology, we cannot even define the Hodge–Riemann relations.

While there does not appear to be a way to salvage the Poincaré duality and Hodge–Riemann relations aspects of the Kähler package, there are weaker Lefschetz properties which the singular cohomology ring may satisfy. We show in Section 5 that the singular cohomology ring of a uniform matroid satisfies what we call the *quasi-projective strong Lefschetz property*.

Theorem 1.6. *Let $\ell \in \text{Gr}_2^W H^2(X_{\Sigma_{U_{r,n}}})$ be generic. Then the map*

$$\times \ell^d : \text{Gr}_j^W H^k(X_{\Sigma_{U_{r,n}}}) \rightarrow \text{Gr}_{j+2d}^W H^{k+2d}(X_{\Sigma_{U_{r,n}}})$$

is injective for $0 \leq d \leq r + j - 2k - 1$ and surjective for $d > r + j - 2k - 1$.

Our proof uses the construction of our basis for the singular cohomology ring to deduce this Lefschetz property from the Hard Lefschetz property of the permutohedral variety.

This strong Lefschetz property also has an interesting combinatorial implication: Each refined Hodge–Poincaré polynomial $\underline{H}_{U_{r,n}}^i(x)$ has unimodal coefficients. In fact, using the formulas from Theorem 1.4, we have verified that the polynomials $\underline{H}_{U_{r,n}}^i(x)$ have ultra-log-concave coefficients for $n \leq 230$. This suggests these polynomials may be real-rooted.

Acknowledgements. This work was supported by NSF RTG grants DMS-2231492 and DMS-2231565. We thank Christin Bibby and Eric Katz for helpful conversations.

2. BACKGROUND AND NOTATION

We assume that the reader is familiar with matroids and toric varieties, and we mostly set notation here. We refer to [Ox11] and [CLS11], respectively, for further details.

2.1. Matroids and Bergman fans. Let M be a loopless matroid of rank r on ground set $[n]$. We write $\mathcal{L}(M)$ for the *lattice of flats* of M . A *proper flat* is flat of rank at most $r - 1$, and a *hyperplane* is flat of rank exactly $r - 1$. We write $\mathcal{H}(M)$ for the set of hyperplanes of M . For a flat $F \in \mathcal{L}(M)$, we write the *restriction of M to F* as M^F .

Throughout this paper we will mostly be restricting ourselves to uniform matroids. For integers $1 \leq r \leq n$, we write $U_{r,n}$ for the uniform matroid of rank r on ground set $[n]$.


Associated to M is a rational fan in $\mathbb{R}^n / \mathbb{R} \cdot (1, \dots, 1)$ called the *Bergman fan*, $\Sigma_{U_{r,n}}$. To construct it, let N be the lattice $\mathbb{Z}^n / \mathbb{Z} \cdot (1, \dots, 1)$, and let e_i be the image of the i th basis vector of \mathbb{Z}^n in N . For a proper, non-empty flat $F \in \mathcal{L}(M)$, define $e_F = \sum_{i \in F} e_i$. The rays of Σ_M are given by $\rho_F = \text{pos}(e_F)$ for each proper, non-empty flat F . The higher-dimensional cones in Σ_M are given by chains of flats in $\mathcal{L}(M)$: $\text{pos}(\rho_{F_1}, \dots, \rho_{F_s}) \in \Sigma_M$ if and only if (up to reordering) $F_1 < \dots < F_s$. Figure 1 depicts $\Sigma_{U_{3,3}}$.

2.2. Singular cohomology of smooth toric varieties. Let N be a lattice and $N_{\mathbb{R}} = N \otimes \mathbb{R}$. Given a unimodular fan $\Sigma \subseteq N_{\mathbb{R}}$, we write X_{Σ} for the associated smooth toric variety defined over \mathbb{C} . We denote the character lattice of X_{Σ} by N^{\vee} and define $N_{\mathbb{Q}}^{\vee} = N^{\vee} \otimes \mathbb{Q}$.

The singular cohomology of X_{Σ} (with rational coefficients) has a description in terms of the Stanley–Reisner ring of Σ and Tor algebras.

Definition 2.1. The *Stanley–Reisner ring* of Σ is the quotient ring

$$\mathbb{Q}[\Sigma] = \frac{\mathbb{Q}[x_{\rho} : \rho \text{ a ray in } \Sigma]}{(x_{\rho_1} \cdots x_{\rho_s} : \text{pos}(\rho_1, \dots, \rho_s) \notin \Sigma)}.$$

FIGURE 1. The Bergman fan of $U_{3,3}$.

The rational embedding $\Sigma \subseteq N_{\mathbb{R}}$ endows $\mathbb{Q}[\Sigma]$ with the structure of a $\text{Sym}(N_{\mathbb{Q}}^{\vee})$ -algebra via the map

$$N^{\vee} \longrightarrow \mathbb{Q}[\Sigma] \quad m \longmapsto \sum_{\rho \text{ ray}} \langle m, u_{\rho} \rangle \cdot x_{\rho},$$

where $\langle -, - \rangle$ is the pairing $N^{\vee} \otimes N$ and $u_{\rho} \in N$ is the primitive generator of the ray ρ .

Theorem 2.2 ([FF24]). *If X_{Σ} is a smooth toric variety, there is a natural ring isomorphism*

$$H^{\bullet}(X_{\Sigma}, \mathbb{Q}) \cong \text{Tor}_{\bullet}^{\text{Sym}(N_{\mathbb{Q}}^{\vee})}(\mathbb{Q}[\Sigma], \mathbb{Q})_{\bullet},$$

where \mathbb{Q} is the residue field of the maximal homogeneous ideal.

The singular cohomology has the additional structure of the *weight filtration*

$$0 = W_{i-1}H^i(X_{\Sigma}) \subseteq W_iH^i(X_{\Sigma}) \subseteq \cdots \subseteq W_{2i-1}H^i(X_{\Sigma}) \subseteq W_{2i}H^i(X_{\Sigma}) = H^i(X_{\Sigma}).$$

By [Tot14], the weight filtration splits, and by [Web04], the associated graded correspond to the bigrading of the Tor algebra

$$\text{Gr}_{2j}^W H^{2j-i}(X_{\Sigma}) \cong \text{Tor}_i^{\text{Sym}(N_{\mathbb{Q}}^{\vee})}(\mathbb{Q}[\Sigma], \mathbb{Q})_j.$$

We call the dimensions of the associated graded *Hodge numbers*.

One obtains a combinatorial description of the singular cohomology by computing the Tor algebra from the Koszul resolution of \mathbb{Q} . Let $\mathbb{Q}[\Sigma]_i$ denote the space of degree i homogeneous forms.

Definition 2.3. The *Koszul complex* of $\mathbb{Q}[\Sigma]$ is the bigraded differential algebra

$$K_i(\mathbb{Q}[\Sigma])_j = \mathbb{Q}[\Sigma]_{j-i} \otimes_{\mathbb{Q}} \bigwedge^i N_{\mathbb{Q}}^{\vee}$$

with differential d defined by the Leibniz rule, $d(x_{\rho}) = 0$, and $d(m) = \sum_{\rho \text{ ray}} \langle m, u_{\rho} \rangle \cdot x_{\rho}$. The algebra structure is induced from the algebra structure on $\mathbb{Q}[\Sigma]$ and $\bigwedge^{\bullet} N_{\mathbb{Q}}^{\vee}$.

Taking homology of the Koszul complex yields *Koszul homology* $H_{\bullet}(K(\mathbb{Q}[\Sigma]))_{\bullet}$, and we have isomorphisms of graded rings

$$\text{Gr}_{2j}^W H^{2j-i}(X_{\Sigma}) \cong \text{Tor}_i^{\text{Sym}(N_{\mathbb{Q}}^{\vee})}(\mathbb{Q}[\Sigma], \mathbb{Q})_j \cong H_i(K(\mathbb{Q}[\Sigma]))_j.$$

2.3. Singular cohomology of matroids. The main object of our paper is the following.

Definition 2.4. For a loopless matroid M , the *singular cohomology ring* of M is the singular cohomology ring of the toric variety X_{Σ_M} .

We use Koszul homology to work with these cohomology rings, and we adopt some notation for our purposes.

The indeterminates of $\mathbb{Q}[\Sigma_M]$ are in bijection with the proper, non-empty flats in $\mathcal{L}(M)$. For a flat F we write x_F for the corresponding indeterminate. The square-free monomials in $\mathbb{Q}[\Sigma_M]$ are in bijection with chains of proper, non-empty flats in $\mathcal{L}(M)$. For such a chain \mathcal{F} , we define $x_{\mathcal{F}} = \prod_{F \in \mathcal{F}} x_F$. Often we will want to consider chains containing the empty flat, so by convention we define $x_{\emptyset} = 1$ and extend the definition of $x_{\mathcal{F}}$ to chains of proper flats.

For Σ_M , $N_{\mathbb{Q}}^{\vee}$ is spanned by elements $\ell_i - \ell_j$ for $i, j \in [n]$, where the ℓ_i form a dual basis to the canonical vectors $e_i \in \mathbb{Z}^n$.

We now give some explicit examples of computations in the Koszul complex.

Example 2.5. For $U_{1,n}$, $\mathbb{Q}[\Sigma_{U_{1,n}}] \cong \mathbb{Q}$. Therefore $K_{\bullet}(\mathbb{Q}[\Sigma_{U_{1,n}}]) \cong \wedge^{\bullet} N_{\mathbb{Q}}^{\vee}$ with trivial differential, and $H_{\bullet}(\mathbb{Q}[\Sigma_{U_{1,n}}]) \cong \wedge^{\bullet} N_{\mathbb{Q}}^{\vee}$.

Example 2.6. For $U_{2,3}$, $\mathbb{Q}[\Sigma_{U_{2,3}}] \cong \frac{\mathbb{Q}[x_1, x_2, x_3]}{x_1 x_2, x_1 x_3, x_2 x_3}$. By [Bin26, Lemma A.5], we can restrict to the quasi-isomorphic *square-free Koszul complex* with generators

$$1, x_1, x_2, x_3, (\ell_1 - \ell_2), (\ell_2 - \ell_3), (\ell_1 - \ell_2) \wedge (\ell_2 - \ell_3), x_1 \otimes (\ell_2 - \ell_3), x_2 \otimes (\ell_1 - \ell_3), x_3 \otimes (\ell_1 - \ell_2).$$

The only non-zero differentials are $d(\ell_1 - \ell_2) = x_1 - x_2$, $d(\ell_2 - \ell_3) = x_2 - x_3$, and $d((\ell_1 - \ell_2) \wedge (\ell_2 - \ell_3)) = x_1 \otimes (\ell_2 - \ell_3) - x_2 \otimes (\ell_1 - \ell_3) + x_3 \otimes (\ell_1 - \ell_2)$.

Therefore, $H_{\bullet}(\mathbb{Q}[\Sigma_{U_{2,3}}])$ has a basis $1, x_3, x_2 \otimes (\ell_1 - \ell_3), x_3 \otimes (\ell_1 - \ell_2)$.

Example 2.7. Take the element $\alpha = x_2 x_{235} \otimes (\ell_3 - \ell_5) \wedge (\ell_1 - \ell_6) \in K_2(\mathbb{Q}[\Sigma_{U_{5,6}}])_4$. We compute its differential to be

$$d(\alpha) = x_2 x_{23} x_{235} \otimes (\ell_1 - \ell_6) - x_2 x_{25} x_{235} \otimes (\ell_1 - \ell_6) - x_2 x_{235} x_{1235} \otimes (\ell_3 - \ell_5) + x_2 x_{235} x_{2356} \otimes (\ell_3 - \ell_5).$$

On the other hand, the chain $\beta = x_3 x_{1234} \otimes (\ell_5 - \ell_6) \in K_1(\mathbb{Q}[\Sigma_{U_{5,6}}])_3$ is a cycle.

3. A BASIS FOR THE SINGULAR COHOMOLOGY RING OF A UNIFORM MATROID

In this section we construct an explicit, combinatorial basis for the singular cohomology ring of a uniform matroid in terms of Koszul homology $H_{\bullet}(K(\mathbb{Q}[\Sigma_{U_{r,n}}]))$.

3.1. Retral and weakly retral chains. We describe our basis for the singular cohomology ring in terms of *retral* and *weakly retral* chains of flats in $\mathcal{L}(U_{r,n})$ which we define here. Roughly speaking, the retral property is a maximality condition which is enforced on each covering relation appearing in the chain. The weakly retral condition is the same, only it is *not* enforced for hyperplanes covering another flat.

Definition 3.1. For proper flats F, F' in $\mathcal{L}(U_{r,n})$, we say that F' *covers* F if $F < F'$ and there is no $F'' \in \mathcal{L}(U_{r,n})$ with $F < F'' < F'$. Moreover, F' is the *maximal cover* of F if F' is the lexicographically largest flat which covers F . As $F' \setminus F$ is a singleton whenever a proper flat F' covers F , this last condition is equivalent to $\max([n] \setminus F) = F' \setminus F$.

For notation, we will always assume that a chain of proper flats in $\mathcal{L}(U_{r,n})$ contains the empty flat. We also write $\max(\mathcal{F})$ for the maximal element in \mathcal{F} .

Definition 3.2. Let $\mathcal{F} = \{\emptyset = F_0 < F_1 < \cdots < F_s\}$ be a chain of proper flats in $\mathcal{L}(U_{r,n})$, and extend notation so that $F_{s+1} = [n]$. We say that \mathcal{F} is *retral* in $\mathcal{L}(U_{r,n})$ if, for all $1 \leq i \leq s$, F_i covers F_{i-1} if and only if F_i is the maximal cover of F_{i-1} in $\mathcal{L}(U_{r,n}^{F_{i+1}})$.

Example 3.3. Below is the set of all retral chains in $\mathcal{L}(U_{3,4})$.

$$\begin{aligned} & \emptyset, \\ & \emptyset < 4, \emptyset < 12, \emptyset < 13, \emptyset < 14, \emptyset < 23, \emptyset < 24, \emptyset < 34, \\ & \emptyset < 4 < 34 \end{aligned}$$

We note that all chains $\emptyset < F$ for $\text{rk}(F) = 2$ are retral as there are no covering relations in the chain; therefore, the maximal cover condition is vacuous.

Example 3.4. The chain $\emptyset < 3 < 123 < 1235$ is a retral chain in $\mathcal{L}(U_{5,5})$. Note, however, that 3 is not the maximal cover of \emptyset in $\mathcal{L}(U_{5,5})$. The maximal cover condition only requires that 3 be the maximal cover of \emptyset in $\mathcal{L}(U_{5,5}^{123})$.

For a chain of proper flats containing a hyperplane, we will remove the maximal cover condition on the hyperplane to define *weakly retral chains*.

Definition 3.5. Let $\mathcal{F} = \{\emptyset = F_0 < F_1 < \cdots < F_s\}$ be a chain of proper flats in $\mathcal{L}(U_{r,n})$ such that $\max(\mathcal{F}) = F_s$ is a hyperplane. We say that \mathcal{F} is *weakly retral* in $\mathcal{L}(U_{r,n})$ if, for all $1 \leq i < s$, F_i covers F_{i-1} if and only if F_i is the maximal cover of F_{i-1} in $\mathcal{L}(U_{r,n}^{F_{i+1}})$.

Example 3.6. The weakly retral chains in $\mathcal{L}(U_{3,4})$ are

$$\begin{aligned} & \emptyset < 12, \emptyset < 13, \emptyset < 14, \emptyset < 23, \emptyset < 24, \emptyset < 34, \\ & \emptyset < 2 < 12, \emptyset < 3 < 13, \emptyset < 4 < 14, \emptyset < 3 < 23, \emptyset < 4 < 24, \emptyset < 4 < 34. \end{aligned}$$

Note that, even though $\emptyset < 4$ is a retral chain, it is not weakly retral as it does not contain a hyperplane.

For a proper, non-empty flat F , we define retral and weakly retral chains of flats in $\mathcal{L}(U_{r,n}^F)$.

Definition 3.7. A chain of proper flats \mathcal{F} in $\mathcal{L}(U_{r,n}^F)$ is *retral* (*weakly retral*) if and only if it is the image of a retral (weakly retral) chain of $\mathcal{L}(U_{|F|,|F|})$ under the unique lexicographic-order preserving isomorphism $U_{|F|,|F|} \rightarrow U_{r,n}^F$.

We need one final definition before describing our basis for the singular cohomology ring.

Definition 3.8. Let \mathcal{F} be a weakly retral chain in $\mathcal{L}(U_{r,n})$ containing the hyperplane $\max(\mathcal{F})$. A pure wedge $\xi = (\ell_{u_1} - \ell_{v_1}) \wedge \cdots \wedge (\ell_{u_i} - \ell_{v_i}) \in \wedge^{i>0} N_{\mathbb{Q}}^{\vee}$ is *admissible* to \mathcal{F} if:

- (1) Each u_m and v_m is in $[n] \setminus \max(\mathcal{F})$.
- (2) $u_1 < \cdots < u_i$, and for all m , v_m is the successor of u_m in the ordered set $[n] \setminus \max(\mathcal{F})$.
- (3) If $\max(\mathcal{F})$ covers $F \in \mathcal{F}$, then u_1 is less than the single element of $\max(\mathcal{F}) \setminus F$.

We note that condition (1) ensures that $x_{\mathcal{F}} \otimes \xi$ is a cycle in the Koszul complex, and condition (2) ensures the collection of admissible wedges to \mathcal{F} are linearly independent. Only condition (3) has non-trivial content for describing our basis.

Example 3.9. For the weakly retral chain $\mathcal{F} = \{\emptyset < 4 < 24\}$ in $\mathcal{L}(U_{3,4})$, the only admissible wedge is $(\ell_1 - \ell_3)$. On the other hand, the weakly retral chain $\mathcal{F} = \{\emptyset < 4 < 14\}$ has no admissible wedges.

Example 3.10. If \mathcal{F} is a weakly retral chain in $\mathcal{L}(U_{n,n})$, then there are no admissible wedges. This is because $[n] \setminus \max(\mathcal{F})$ is a singleton, so conditions (1) and (2) cannot be simultaneously satisfied.

3.2. A basis of the singular cohomology ring. We now describe and construct our basis for the singular cohomology ring of a uniform matroid.

Theorem 3.11. *The set*

$$\{x_{\mathcal{F}} : \mathcal{F} \text{ is retral}\} \cup \{x_{\mathcal{F}} \otimes \xi : \mathcal{F} \text{ is weakly retral and } \xi \text{ is admissible to } \mathcal{F}\}$$

is a basis of $H_{\bullet}(K(\mathbb{Q}[\Sigma_{U_{r,n}}])) \cong H^{\bullet}(X_{\Sigma_{U_{r,n}}})$ as a \mathbb{Q} -vector space.

We will prove Theorem 3.11 through the following five lemmas, inducting on r . We start with the base case.

Lemma 3.12. *Theorem 3.11 holds in rank 1.*

Proof. From Example 2.5, $H_{\bullet}(K(\mathbb{Q}[\Sigma_{U_{1,n}}])) \cong \wedge^{\bullet} N_{\mathbb{Q}}^{\vee}$. The chain \emptyset is the only weakly retral chain (it is also retral), and the wedges which are admissible to \emptyset form a basis of $\wedge^{>0} N_{\mathbb{Q}}^{\vee}$. \square

For the inductive step of the proof, we use the *truncation exact sequence*

$$(2) \quad \cdots \rightarrow \bigoplus_{H \in \mathcal{H}(U_{r,n})} H_i(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_j \rightarrow H_i(K(\mathbb{Q}[\Sigma_{U_{r,n}}]))_{j+1} \rightarrow H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))_{j+1} \rightarrow \cdots$$

where the subfan $\overline{\text{St}}(\rho_H) = \{\sigma \in \Sigma_{U_{r,n}} : \text{pos}(\sigma, \rho_H) \in \Sigma_{U_{r,n}}\}$ is the *closed star* of ρ_H . The exact sequence is induced by applying Koszul homology to the exact sequence of $\text{Sym}(N_{\mathbb{Q}}^{\vee})$ -algebras

$$0 \rightarrow \bigoplus_{H \in \mathcal{H}(U_{r,n})} \mathbb{Q}[\overline{\text{St}}(\rho_H)] \xrightarrow{\oplus \cdot x_{\rho_H}} \mathbb{Q}[\Sigma_{U_{r,n}}] \rightarrow \mathbb{Q}[\Sigma_{U_{r-1,n}}] \rightarrow 0.$$

We will write the connecting homomorphism as

$$\delta_i : H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))_j \rightarrow \bigoplus_H H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_{j-1}.$$

The inductive hypothesis provides bases for $H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))_j$ and each $H_i(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_j$ which we describe in Lemma 3.14. In Lemmas 3.16 and 3.17, we use these bases to compute a Gröbner basis for the image of the connecting homomorphism δ_i . We describe the standard monomials for $\text{coker } \delta_i$ in Lemma 3.18, and these produce the basis for Theorem 3.11.

Definition 3.13. For a hyperplane $H \in \mathcal{H}(U_{r,n})$, define $P_H \subseteq \wedge^{>0} N_{\mathbb{Q}}^{\vee}$ to be the set of pure wedges $\xi = (\ell_{u_1} - \ell_{v_1}) \wedge \cdots \wedge (\ell_{u_i} - \ell_{v_i})$ such that

- (1) Each u_m and v_m is in $[n] \setminus H$.
- (2) $u_1 < \cdots < u_i$, and for all m , v_m is the successor of u_m in the ordered set $[n] \setminus H$.

Lemma 3.14. *Assume that Theorem 3.11 holds in rank $r - 1$. Then*

$$\{x_{\mathcal{F}} \otimes \xi : \mathcal{F} \text{ is retral in } \mathcal{L}(U_{r,n}^H) \text{ and } \xi \in P_H\}$$

is a basis for $H_{\bullet}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$, where $H \in \mathcal{H}(U_{r,n})$.

Proof. By [Bin26, Lemma A.3], $H_{\bullet}(K(\mathbb{Q}[\text{St}(\rho_H)])) \cong H_{\bullet}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$, where $\text{St}(\rho_H)$ is the *star* of ρ_H in $\Sigma_{U_{r,n}}$ (see [CLS11, Definition (3.2.8)]). This isomorphism is induced by the natural inclusion of complexes $K_{\bullet}(\mathbb{Q}[\text{St}(\rho_H)]) \hookrightarrow K_{\bullet}(\mathbb{Q}[\overline{\text{St}}(\rho_H)])$. In our case, $\text{St}(\rho_H) \cong \Sigma_{U_{r-1,r-1}} \times \emptyset$ in the product vector space $N_{H,\mathbb{R}} \times N_{[n] \setminus H,\mathbb{R}}$ where $N_{H,\mathbb{R}} = \text{span}(e_i : i \in H)/e_H$ and $N_{[n] \setminus H,\mathbb{R}} = \text{span}(e_i : i \in [n] \setminus H)/e_{[n] \setminus H}$. Therefore, $K_{\bullet}(\mathbb{Q}[\text{St}(\rho_H)])$ is isomorphic to the tensor complex $K_{\bullet}(\mathbb{Q}[\Sigma_{U_{r-1,r-1}}]) \otimes_{\mathbb{Q}} \wedge^{\bullet} N_{[n] \setminus H,\mathbb{Q}}^{\vee}$, with trivial differential on $\wedge^{\bullet} N_{[n] \setminus H,\mathbb{Q}}^{\vee}$. As P_H forms a basis

for $N_{[n] \setminus H, \mathbb{Q}}^\vee$, the lemma follows from the Künneth formula and the basis for $H_\bullet(K(\mathbb{Q}[\Sigma_{U_{r-1}, r-1}])))$ described in Theorem 3.11. \square

For each $H \in \mathcal{H}(U_{r,n})$, choose an arbitrary ordering of the basis elements from Lemma 3.14. Then extend this to an ordering of the basis elements of $\bigoplus_H H_\bullet(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])))$ such that a basis element of $H_\bullet(K(\mathbb{Q}[\overline{\text{St}}(\rho_{H_1}])))$ precedes a basis element of $H_\bullet(K(\mathbb{Q}[\overline{\text{St}}(\rho_{H_2}])))$ if and only if $H_1 <_{\text{lex}} H_2$.

We now study the connecting homomorphisms δ_i through Gröbner theory. For a basis element α of $H_i(K(\mathbb{Q}[\Sigma_{U_{r-1}, n}])))$, we are interested in the first basis element of $\bigoplus_H H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])))$ that appears in $\delta_i(\alpha)$.

Definition 3.15. For a basis element $\alpha = x_{\mathcal{F}} \otimes \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m}) \in H_i(K(\mathbb{Q}[\Sigma_{U_{r-1}, n}])))$, define

$$\text{in}(\alpha) = x_{\mathcal{F}} \otimes \bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m}) \in H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))),$$

where $H = \max(\mathcal{F}) \cup u_1$ is a hyperplane of $U_{r,n}$.

Lemma 3.16. *Assume that Theorem 3.11 holds in rank $r - 1$. Then for each basis element $\alpha \in H_{i>0}(K(\mathbb{Q}[\Sigma_{U_{r-1}, n}])))$, $\text{in}(\alpha)$ is the first basis element appearing in $\delta_i(\alpha)$.*

Proof. We first show that $\text{in}(\alpha)$ is a basis element. Write $\alpha = x_{\mathcal{F}} \otimes \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$ and $H = \max(\mathcal{F}) \cup u_1$ as above. By assumption, \mathcal{F} is a weakly retral chain in $\mathcal{L}(U_{r-1, n})$, and we must show that it is retral in $\mathcal{L}(U_{r,n}^H)$. The only thing to check is the maximal cover condition for $\max(\mathcal{F})$. If $\max(\mathcal{F})$ covers $F \in \mathcal{F}$, then $u_1 < \max(\mathcal{F}) \setminus F$ by the definition of admissible wedges. The two covers of F in $\mathcal{L}(U_{r,n}^H)$ are therefore $F \cup u_1$ and $\max(\mathcal{F})$, and $\max(\mathcal{F})$ is the maximal one. Finally, one immediately verifies that $\bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m}) \in P_H$.

We now show that $\text{in}(\alpha)$ is the first basis element appearing in $\delta_i(\alpha)$. For a hyperplane H_1 , denote the inclusion $\iota_{H_1} : H_\bullet(K(\mathbb{Q}[\overline{\text{St}}(\rho_{H_1}]))) \hookrightarrow \bigoplus_H H_\bullet(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])))$. Then δ_i is given by

$$\delta_i(\alpha) = \sum_{m=1}^i (-1)^{m+1} \left[\iota_{\max(\mathcal{F}) \cup u_m} \left(x_{\mathcal{F}} \otimes \bigwedge_{p \neq m} (\ell_{u_p} - \ell_{v_p}) \right) - \iota_{\max(\mathcal{F}) \cup v_m} \left(x_{\mathcal{F}} \otimes \bigwedge_{p \neq m} (\ell_{u_p} - \ell_{v_p}) \right) \right].$$

By the definition of admissible wedges, $\text{in}(\alpha)$ is the only term in $\delta_i(\alpha)$ involving $\max(\mathcal{F}) \cup u_1$. This is the lexicographically minimal hyperplane involved in $\delta_i(\alpha)$. \square

Lemma 3.17. *Assume that Theorem 3.11 holds in rank $r - 1$. Then $\alpha \mapsto \text{in}(\alpha)$ defines an injective map from basis elements of $H_{i>0}(K(\mathbb{Q}[U_{r-1, n}])))$ to basis elements of $\bigoplus_H H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])))$.*

Proof. If α is a basis element of $H_{i>0}(K(\mathbb{Q}[\Sigma_{U_{r-1}, n}])))$ with $\text{in}(\alpha) = x_{\mathcal{F}} \otimes \bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m})$ in $H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])))$, let $u_1 = H \setminus \max(\mathcal{F})$ and v_1 the successor of u_1 in $[n] \setminus \max(\mathcal{F})$. Then we find that $\alpha = x_{\mathcal{F}} \otimes \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$. \square

A consequence of Lemmas 3.16 and 3.17 is that, for $i > 0$,

$$\{\delta_i(\alpha) : \alpha \text{ is a basis element of } H_i(K(\mathbb{Q}[\Sigma_{U_{r-1}, n}])))\}$$

is a Gröbner basis for $\text{im } \delta_i$.

Lemma 3.18. *Assume that Theorem 3.11 holds in rank $r - 1$. With respect to the above Gröbner basis, the standard monomials for coker δ_i are the following basis elements of $H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$. For $i > 1$,*

$$\bigcup_{H \in \mathcal{H}(U_{r,n})} \left\{ x_{\mathcal{F}} \otimes \bigwedge_{m=1}^{i-1} (\ell_{u_m} - \ell_{v_m}) \in H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])) \mid \begin{array}{l} 1. H \text{ does not cover } \max(\mathcal{F}), \text{ or} \\ 2. u_1 < H \setminus \max(\mathcal{F}) \end{array} \right\}.$$

For $i = 1$,

$$\bigcup_{H \in \mathcal{H}(U_{r,n})} \left\{ x_{\mathcal{F}} \in H_0(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)])) \mid \begin{array}{l} 1. H \text{ does not cover } \max(\mathcal{F}), \text{ or} \\ 2. H \text{ is the maximal cover of } \max(\mathcal{F}) \text{ in } \mathcal{L}(U_{r,n}) \end{array} \right\}.$$

Proof. By definition, the standard monomials are the basis elements of $\bigoplus_H H_{\bullet}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$ which are not in (α) for any basis element $\alpha \in H_{>0}(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))$. We will prove that the basis elements of the form in (α) are precisely those not listed in the statement.

For notation, write $\alpha = x_{\mathcal{F}} \otimes \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$ for a basis element of $H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))$, and let $H = \max(\mathcal{F}) \cup u_1$.

Case $i > 1$: We have $\text{in}(\alpha) = x_{\mathcal{F}} \otimes \bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m}) \in H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$. Then H covers $\max(\mathcal{F})$ and $u_2 \geq H \setminus \max(\mathcal{F})$.

Conversely, suppose we have a basis element $x_{\mathcal{F}} \otimes \bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m})$ in $H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$ with H covering $\max(\mathcal{F})$ and $u_2 \geq H \setminus \max(\mathcal{F})$. Then define $u_1 = H \setminus \max(\mathcal{F})$. We claim that $\alpha = x_{\mathcal{F}} \otimes \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$ is a basis element of $H_{\bullet}(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))$ with $\text{in}(\alpha) = x_{\mathcal{F}} \otimes \bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m})$. To see that it is a basis element, note that $\max(\mathcal{F})$ is a hyperplane of $U_{r-1,n}$, as it is covered by H , a hyperplane of $U_{r,n}$. As \mathcal{F} is retral in $U_{r,n}^H$, \mathcal{F} is weakly retral in $U_{r-1,n}$. The wedge $\bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$ is admissible to \mathcal{F} , as if $\max(\mathcal{F})$ covers a flat F in \mathcal{F} , then $\max(\mathcal{F}) \setminus F > H \setminus \max(\mathcal{F}) = u_1$ because \mathcal{F} is retral in $U_{r,n}^H$. Finally, it is clear that $\text{in}(\alpha) = x_{\mathcal{F}} \otimes \bigwedge_{m=2}^i (\ell_{u_m} - \ell_{v_m})$.

Case $k = 1$: We have $\text{in}(\alpha) = x_{\mathcal{F}} \in H_0(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$. Then H covers $\max(\mathcal{F})$, but it is not the maximal cover of $\max(\mathcal{F})$ in $\mathcal{L}(U_{r,n})$. In particular, $\max(\mathcal{F}) \cup v_1$ is a cover that is lexicographically greater than H .

Conversely, suppose that $x_{\mathcal{F}}$ is a basis element of $H_0(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$ such that H covers $\max(\mathcal{F})$ and H is not the maximal cover of $\max(\mathcal{F})$ in $\mathcal{L}(U_{r,n})$. Let $u_1 = H \setminus \max(\mathcal{F})$, and let v_1 be the successor of u_1 in $[n] \setminus \max(\mathcal{F})$. This successor exists because H is not the maximal cover of \mathcal{F} . We claim that $\alpha = x_{\mathcal{F}} \otimes (\ell_{u_1} - \ell_{v_1})$ is a basis element of $H_1(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))$. It suffices to check that if $\max(\mathcal{F})$ covers F in \mathcal{F} , then $u_1 < \max(\mathcal{F}) \setminus F$. Suppose this is the case. As \mathcal{F} is retral in $\mathcal{L}(U_{r,n}^H)$, $\max(\mathcal{F})$ is the maximal cover of F in $\mathcal{L}(U_{r,n}^H)$. This means that $\max(\mathcal{F}) \setminus F > H \setminus \max(\mathcal{F}) = u_1$. Clearly, $\text{in}(\alpha) = x_{\mathcal{F}} \in H_0(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$. \square

We now deduce our main result.

Proof of Theorem 3.11. Inducting on the rank r , the base case is Lemma 3.12. By Lemma 3.17, the connecting homomorphism $\delta_i: H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}])) \rightarrow \bigoplus_H H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))$ is injective for $i \geq 1$. For $i = 0$, the connecting homomorphism vanishes. Therefore, $H_{\bullet}(K(\mathbb{Q}[\Sigma_{U_{r,n}}])) \cong H_0(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}])) \oplus \bigoplus_{i \geq 1} \text{coker } \delta_i$. To get a basis, we pull back the basis for $H_0(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))$ and push forward the standard monomials from Lemma 3.18. One verifies that the basis elements from the pull-back are

$$\{x_{\mathcal{F}} \mid \mathcal{F} \text{ retral, not containing a hyperplane}\},$$

and the basis elements from the push-forward are

$$\{x_{\mathcal{F}} \mid \mathcal{F} \text{ retral, containing a hyperplane}\} \cup \{x_{\mathcal{F}} \otimes \xi \mid \mathcal{F} \text{ weakly retral and } \xi \text{ admissible to } \mathcal{F}\}.$$

Together, these elements form the desired basis. \square

Corollary 3.19. *The set*

$$\{x_{\mathcal{F}} : \mathcal{F} \text{ retral in } \mathcal{L}(U_{r,n})\}$$

is a basis for the Chow ring $A^\bullet(X_{\Sigma_{U_{r,n}}})$ as a \mathbb{Q} -vector space.

Example 3.20. The Betti diagram for the singular cohomology of $U_{3,4}$ is

	0	1
0	1	0
1	7	6
2	1	3

The basis we have given for $H_\bullet(K(\mathbb{Q}[\Sigma_{U_{3,4}}]))$ is

	0	1
0	1	-
1	$x_4, x_{12}, x_{13}, x_{14}, x_{23}, x_{24}, x_{34}$	$x_{12} \otimes (\ell_3 - \ell_4), x_{13} \otimes (\ell_2 - \ell_4), x_{14} \otimes (\ell_2 - \ell_3),$ $x_{23} \otimes (\ell_1 - \ell_4), x_{24} \otimes (\ell_1 - \ell_3), x_{34} \otimes (\ell_1 - \ell_2)$
2	$x_4 x_{34}$	$x_3 x_{23} \otimes (\ell_1 - \ell_4), x_4 x_{24} \otimes (\ell_1 - \ell_3), x_4 x_{34} \otimes (\ell_1 - \ell_2)$

arranged according to this Betti diagram.

4. COMBINATORICS OF RETRAL CHAINS

In this section, we study the combinatorics of retral and weakly retral chains in $\mathcal{L}(U_{r,n})$. From this we produce multiple formulas for the refined Hodge–Poincaré polynomials. When restricted to the special case of Chow polynomials, these formulas give combinatorial proofs for results of Hoster [Hos26], Hameister, Rao, and Simpson [HRS21], and Ferroni et al. [Fer+24].

4.1. Rank-selected retral chains. We first count retral chains whose flats are of specified rank. This produces a combinatorial proof of a formula for the Chow polynomial of a uniform matroid due to Hoster [Hos26].

Definition 4.1. Let \mathcal{F} be a chain of proper flats of $\mathcal{L}(U_{r,n})$. We define the *ranks* of \mathcal{F} to be the subset $\text{rks}(\mathcal{F}) = \{j \in \{0, \dots, r-1\} : \mathcal{F} \text{ contains a flat of rank } j\}$. For $i \in \text{rks}(\mathcal{F})$, we write F^i for the flat in \mathcal{F} of rank i .

Recall our convention that every chain \mathcal{F} contains the empty flat.

Definition 4.2. Let $R \subseteq \{0, \dots, r-1\}$ be a subset containing 0. Define $C_R(U_{r,n})$ to be the set of retral chains \mathcal{F} in $\mathcal{L}(U_{r,n})$ such that $\text{rks}(\mathcal{F}) = R$.

For $R \subseteq \{0, \dots, r-1\}$, partition $R = R_1 \sqcup \dots \sqcup R_s$ into its maximal consecutive subsets, ordered so that $\min R_i < \min R_{i+1}$. Define the multinomial coefficient

$$\binom{n}{\Delta R} = \binom{n}{\min R_1 - 0, \min R_2 - \min R_1, \dots, \min R_s - \min R_{s-1}}$$

and the index $\lfloor R \rfloor = \min R_s$.

One way to construct a retral chain $\mathcal{F} \in C_R(U_{r,n})$ is to choose an arbitrary chain of flats \mathcal{G} with $\text{rks}(\mathcal{G}) = \{\min R_1, \dots, \min R_s\}$ and then uniquely extend using the retral condition.

Lemma 4.3. *Let R be a subset of $\{0, \dots, r-1\}$ which contains 0, and partition $R = R_1 \sqcup \dots \sqcup R_s$ as above. Given a chain of flats $\mathcal{G} \subseteq \mathcal{L}(U_{r,n})$ with $\text{rks}(\mathcal{G}) = \{\min R_1, \dots, \min R_s\}$, there is a unique $\mathcal{F} \in C_R(U_{r,n})$ which extends \mathcal{G} .*

Proof. A chain of proper flats \mathcal{F} with $\text{rks}(\mathcal{F}) = R$ is retral if and only if each sequence of integers $(F^{\min R_{j+1}} \setminus F^{\min R_j}, \dots, F^{\max R_j} \setminus F^{\max R_{j-1}})$ is decreasing and $F^{\max R_j} \setminus F^{\max R_{j-1}}$ equals the maximal element of $F^{\min R_{j+1}} \setminus F^{\max R_{j-1}}$ (here we use the convention that $[n] = F^{\min R_{s+1}}$). This is satisfied if and only if $(F^{\min R_{j+1}} \setminus F^{\min R_j}, \dots, F^{\max R_j} \setminus F^{\max R_{j-1}})$ is the sequence of the $(\max R_j - \min R_j)$ largest elements of $F^{\min R_{j+1}} \setminus F^{\min R_j}$ written in decreasing order. Therefore, there is exactly one such $\mathcal{F} \in C_R(U_{r,n})$ which extends \mathcal{G} , as \mathcal{G} determines each $F^{\min R_i}$. \square

The number of chains $\mathcal{G} \subseteq \mathcal{L}(U_{r,n})$ with $\text{rks}(\mathcal{G}) = \{\min R_1, \dots, \min R_s\}$ is given by $\binom{n}{\Delta R}$. As a consequence, we have the following count of rank-selected retral chains.

Proposition 4.4. *Let R be a subset of $\{0, \dots, r-1\}$ which contains 0. Then $|C_R(U_{r,n})| = \binom{n}{\Delta R}$.*

The basis of Theorem 3.11 thus gives a combinatorial proof, in terms of retral chains of flats, of the following result of Hoster [Hos26, Theorem 3]³.

Theorem 4.5. *For $r \geq 1$,*

$$\underline{H}_{U_{r,n}}(x) = \sum_{\substack{R \subseteq \{0, \dots, r-1\} \\ 0 \in R}} \binom{n}{\Delta R} \cdot x^{|R|-1}.$$

4.2. Rank-selected weakly retral chains and admissible wedges. Our next goal is count the other basis elements $x_{\mathcal{F}} \otimes \xi$, where \mathcal{F} is a weakly retral chain in $\mathcal{L}(U_{r,n})$ with prescribed ranks and ξ is admissible to \mathcal{F} .

Definition 4.6. Let R be a subset of $\{0, \dots, r-1\}$ which contains 0 and $r-1$. Define $W_{R,i}(U_{r,n})$ to be the set of pairs (\mathcal{F}, ξ) such that \mathcal{F} is a weakly retral chain in $\mathcal{L}(U_{r,n})$, $\text{rks}(\mathcal{F}) = R$, and $\xi \in \wedge^i N_{\mathbb{Q}}^{\vee}$ is admissible to \mathcal{F} .

Our method of constructing pairs $(\mathcal{F}, \xi) \in W_{R,i}(U_{r,n})$ is similar to our construction of elements of $C_R(U_{r,n})$ in that we will begin with an arbitrary chain \mathcal{G} with $\text{rks}(\mathcal{G}) = \{\min R_1, \dots, \min R_s\}$. However, this is not enough to uniquely determine (\mathcal{F}, ξ) from \mathcal{G} , as there is some freedom in the choice of hyperplane of \mathcal{F} , and \mathcal{G} imposes few restrictions on $\xi = \wedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$. Our method of determining elements $(\mathcal{F}, \xi) \in W_{R,i}(U_{r,n})$ is outlined as follows:

- (1) Choose an arbitrary chain \mathcal{G} with $\text{rks}(\mathcal{G}) = \{\min R_1, \dots, \min R_s\}$,
- (2) choose u_1 ,
- (3) choose the hyperplane of \mathcal{F} ,
- (4) choose the remaining u_2, \dots, u_i ,

then extend to a unique pair (\mathcal{F}, ξ) . The choice in (1) will be arbitrary, but there will be some restrictions in (2)–(4).

Lemma 4.7. *Let R be a subset of $\{0, \dots, r-1\}$ which contains 0 and $r-1$. Partition $R = R_1 \sqcup \dots \sqcup R_s$ as above. Given*

- (1) *a chain of flats $\mathcal{G} \subseteq \mathcal{L}(U_{r,n})$ with $\text{rks}(\mathcal{G}) = \{\min R_1, \dots, \min R_s\}$,*
- (2) *an integer $1 \leq \ell \leq n - r - i + 1$,*
- (3) *an $(r - \lfloor R \rfloor - 1)$ element subset $F \subseteq [n] \setminus G^{\lfloor R \rfloor}$ with $\min F$ greater than the ℓ th smallest element of $[n] \setminus G^{\lfloor R \rfloor}$,*

³Our indices differ by a shift from Hoster's.

(4) an $(i - 1)$ -element subset $I \subseteq [n] \setminus (G^{\lfloor R \rfloor} \cup F)$ with $\min I$ greater than the ℓ th smallest element of $[n] \setminus G^{\lfloor R \rfloor}$ and $\max I < \max [n] \setminus (G^{\lfloor R \rfloor} \cup F)$,

there is a unique pair $(\mathcal{F}, \xi = \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})) \in W_{R,i}(U_{r,n})$ such that

- (a) $\max(\mathcal{F}) = F \cup G^{\lfloor R \rfloor}$,
- (b) \mathcal{F} extends \mathcal{G} ,
- (c) u_1 is the ℓ th smallest element of $[n] \setminus G^{\lfloor R \rfloor}$,
- (d) $I = \{u_2, \dots, u_i\}$.

Proof. Similar to the proof of Lemma 4.3, there is a unique weakly retral chain \mathcal{F} with $\text{rks}(\mathcal{F}) = R$ satisfying conditions (a) and (b). Conditions (c) and (d) uniquely determine $\xi = \bigwedge_{m=1}^i (\ell_{u_m} - \ell_{v_m})$, as we must have $u_1 < u_2 < \dots < u_i$ and v_m must be the successor of u_m in $[n] \setminus \max(\mathcal{F})$. Note that the successor of u_i exists because of the restriction in (4).

It remains to see that ξ is admissible to \mathcal{F} . Indeed, if F^{r-1} covers F^{r-2} in \mathcal{F} , then $u_1 < F^{r-1} \setminus F^{r-2}$ due to the restriction in (3) and condition (c). \square

It is straightforward to see that every pair $(\mathcal{F}, \xi) \in W_{R,i}(U_{r,n})$ arises from Lemma 4.7 for a unique set of data \mathcal{G}, ℓ, F, I . The following counts the possible choices of \mathcal{G}, ℓ, F , and I above.

Proposition 4.8. *Let R be a subset of $\{0, \dots, r - 1\}$ containing 0 and $r - 1$. Then*

$$|W_R(U_{r,n})| = \binom{n}{\Delta R} \sum_{\ell=1}^{n-r-i+1} \binom{n - \lfloor R \rfloor - \ell}{r - \lfloor R \rfloor - 1} \binom{n - r - \ell}{i - 1}.$$

Summing over appropriate subsets R produces the refined Hodge–Poincaré polynomials.

Theorem 4.9. *For $1 \leq i \leq n - r$,*

$$\underline{H}_{U_{r,n}}^i(x) = \sum_{\substack{R \subseteq \{0, \dots, r-1\} \\ 0, r-1 \in R}} \sum_{\ell=1}^{n-r-i+1} \binom{n}{\Delta R} \binom{n - \lfloor R \rfloor - \ell}{r - \lfloor R \rfloor - 1} \binom{n - r - \ell}{i - 1} \cdot x^{|R|-1}.$$

4.3. Refined Hodge–Poincaré polynomials and derangement polynomials. In this subsection, we give another way of constructing retral chains, this time through concatenation. This leads to combinatorial proofs of results from [HRS21] and [Fer+24] which express the Chow polynomial $\underline{H}_{U_{r,n}}(x)$ in terms of the *derangement polynomials*. We then obtain similar formulas for the other refined Hodge–Poincaré polynomials.

Definition 4.10. Let \mathfrak{D}_n denote the set of *derangements* on $[n]$ —these are the permutations in \mathfrak{S}_n with no fixed points. Recall that an *excedence* of $\sigma \in \mathfrak{S}_n$ is an index $i \in [n]$ such that $i < \sigma(i)$. We write $\text{exc}(\sigma)$ for the number of excedences of σ .

For $n \geq 1$, the n th *derangement polynomial* is

$$d_n(x) = \sum_{\sigma \in \mathfrak{D}_n} x^{\text{exc}(\sigma)},$$

with $d_0(x) = 1$ by convention.

For a chain of flats \mathcal{F} , we define $\mathcal{F}_{\geq j} = \{F^i : i \in \text{rks}(\mathcal{F}) \text{ and } i \geq j\}$ and $\mathcal{F}_{< j}$ similarly.

Lemma 4.11. *Let $j > 1$ and suppose \mathcal{F} is a retral chain in $\mathcal{L}(U_{r,n})$ with $j \in \text{rks}(\mathcal{F})$ but $j - 1 \notin \text{rks}(\mathcal{F})$. Then $\emptyset < \mathcal{F}_{\geq j}$ is a retral chain in $\mathcal{L}(U_{r,n})$ with $\min \text{rks}(\mathcal{F}_{\geq j}) = j$, and $\mathcal{F}_{< j}$ is a retral chain in $\mathcal{L}(U_{r,n}^{F^j})$ with $\max \text{rks}(\mathcal{F}_{< j}) < j - 1$.*

The converse is also true: If $\emptyset < \mathcal{F}_{\geq j}$ is a retral chain in $\mathcal{L}(U_{r,n})$ with $\min \text{rks}(\mathcal{F}_{\geq j}) = j$ and $\mathcal{F}_{< j}$ is a retral chain in $\mathcal{L}(U_{r,n}^{F^j})$ with $\max \text{rks}(\mathcal{F}_{< j}) < j - 1$, then the concatenation $\mathcal{F}_{< j} < \mathcal{F}_{\geq j}$ is a retral chain in $\mathcal{L}(U_{r,n})$.

Proof. The first statement is immediate from the definition of retral chains. The second statement follows because F^j covers no flat in $\mathcal{F}_{< j} < \mathcal{F}_{\geq j}$, so the maximal cover condition on this flat is vacuous. \square

We use this construction to count retral chains in $\mathcal{L}(U_{r,n})$, yielding a formula for $\underline{H}_{U_{r,n}}(x)$. For the following, we use the convention that $\underline{H}_{U_{0,1}}(x) = 0$ and $\underline{H}_{U_{-1,0}}(x) = \frac{1}{x}$.

Proposition 4.12. *For $r \geq 1$,*

$$\underline{H}_{U_{r,n}}(x) = \sum_{j=0}^{r-1} \binom{n}{j} \underline{H}_{U_{j-1,j}}(x) \cdot (x + x^2 + \cdots + x^{r-j}).$$

Proof. For $j \geq 0$, we claim that the generating function, with respect to length, for retral chains \mathcal{F} in $\mathcal{L}(U_{r,n})$ such that j is the maximal index with $j \in \text{rks}(\mathcal{F})$ and $j - 1 \notin \text{rks}(\mathcal{F})$ is

$$\binom{n}{j} \underline{H}_{U_{j-1,j}}(x) \cdot (x + x^2 + \cdots + x^{r-j}).$$

We begin with the edge cases of $j = 0, 1$. For $j = 0$, Lemma 4.3 implies that such retral chains are determined by their length, which ranges from 0 to $r - 1$. The generating function is then $1 + \cdots + x^{r-1}$, which agrees with the claim by our convention for $\underline{H}_{U_{-1,0}}(x)$. For $j = 1$, our convention that all retral chains contain \emptyset implies the generating function is 0 which also agrees with the claim by our convention for $\underline{H}_{U_{0,1}}(x)$.

For $j > 1$, Lemma 4.11 implies that such \mathcal{F} is determined uniquely by $\mathcal{F}_{\geq j}$ and $\mathcal{F}_{< j}$. Note that $\text{rks}(\mathcal{F}_{\geq j})$ is a consecutive sequence of integers, so Lemma 4.3 implies that it is determined by F^j and its length from 1 to $r - j$. On the other hand, $\mathcal{F}_{< j}$ is a retral chain of $\mathcal{L}(U_{r,n}^{F^j})$ with $\max \text{rks}(\mathcal{F}_{< j}) < j - 1$. This is equivalent to a retral chain in $\mathcal{L}(U_{j-1,j})$. Therefore, the generating function is as claimed, and the result follows by summing over j . \square

We now relate the Chow polynomials of uniform matroids to the derangement polynomials.

Proposition 4.13 ([HRS21, Theorem 5.1] and [Fer+24, Theorem 1.9]). *The following formulas for the Chow polynomials hold:*

$$\begin{aligned} x \cdot \underline{H}_{U_{n-1,n}}(x) &= d_n(x) \\ \underline{H}_{U_{r,n}}(x) &= \sum_{j=0}^{r-1} \binom{n}{j} d_j(x) \cdot (1 + x + \cdots + x^{r-1-j}). \end{aligned}$$

Proof. Proposition 4.12 applied to $U_{n-1,n}$ shows that $\{x \cdot \underline{H}_{U_{n-1,n}}(x)\}_{n=0}^{\infty}$ satisfies the recurrence

$$f_n(x) = \sum_{j=0}^{n-2} \binom{n}{j} f_j(x) \cdot (x + \cdots + x^{n-1-j})$$

which the family $\{d_n(x)\}_{n=0}^{\infty}$ also satisfies by [JMS19, Corollary 4.2]. As $x \cdot \underline{H}_{U_{n-1,n}}(x) = d_n(x)$ for $n = 0, 1$, we have $x \cdot \underline{H}_{U_{n-1,n}}(x) = d_n(x)$ for all n .

Proposition 4.12 yields the formula for $\underline{H}_{U_{r,n}}(x)$ after the substitution $d_j(x) = x \cdot \underline{H}_{U_{j-1,j}}(x)$. \square

4.3.1. *Weakly retral chains and admissible wedges.* A similar strategy of concatenation produces a method for constructing pairs (\mathcal{F}, ξ) of a weakly retral chain \mathcal{F} and admissible wedge ξ .

Lemma 4.14. *Let $j > 1$ and suppose that \mathcal{F} is a weakly retral chain in $\mathcal{L}(U_{r,n})$ with $j \in \text{rks}(\mathcal{F})$ but $j - 1 \notin \text{rks}(\mathcal{F})$. Suppose also that ξ is admissible to \mathcal{F} . Then $\emptyset < \mathcal{F}_{\geq j}$ is a weakly retral chain in $\mathcal{L}(U_{r,n})$ with $\min \text{rks}(\mathcal{F}_{\geq j}) = j$, $\mathcal{F}_{< j}$ is a retral chain in $\mathcal{L}(U_{r,n}^{F_j})$ with $\max \text{rks}(\mathcal{F}_{< j}) < j - 1$, and ξ is admissible to $\emptyset < \mathcal{F}_{\geq j}$.*

The converse also holds.

From this construction we derive a formula for the refined Hodge–Poincaré polynomials in terms of the derangement polynomials.

Theorem 4.15. *For $1 \leq i \leq n - r$,*

$$\underline{H}_{U_{r,n}}^i(x) = \sum_{j=0}^{r-1} \sum_{\ell=1}^{n-r-i+1} \binom{n}{j} \binom{n-j-\ell}{r-j-1} \binom{n-r-\ell}{i-1} d_j(x) \cdot x^{r-j-1}.$$

Proof. For $j \geq 0$, we claim the generating function, with respect to length of chain, for pairs $(\mathcal{F}, \xi) \in \bigcup_R W_{R,i}(U_{r,n})$ such that j is the maximal integer with $j \in \text{rks}(\mathcal{F})$ and $j - 1 \notin \text{rks}(\mathcal{F})$ is

$$\sum_{\ell=1}^{n-r-i+1} \binom{n}{j} \binom{n-j-\ell}{r-j-1} \binom{n-r-\ell}{i-1} d_j(x) \cdot x^{r-j-1}.$$

Indeed, Lemma 4.14 implies that such pairs are uniquely determined by $(\emptyset < \mathcal{F}_{\geq j}, \xi) \in W_{R,i}(U_{r,n})$ with $R = \{0, j, j + 1, \dots, r - 1\}$ and a retral chain $\mathcal{F}_{< j} \in \mathcal{L}(U_{r,n}^{F_j})$ with $j - 1 \notin \text{rks}(\mathcal{F}_{< j})$. By Proposition 4.8, the generating function for the former is

$$\sum_{\ell=1}^{n-r-i+1} \binom{n}{j} \binom{n-j-\ell}{r-j-1} \binom{n-r-\ell}{i-1} \cdot x^{r-j},$$

while the generating function for the latter is $d_j(x) \cdot \frac{1}{x}$ by Proposition 4.13. The result follows by summing over j . \square

4.4. Recursive formulas for refined Hodge–Poincaré polynomials. The truncation exact sequence (2) and Lemma 3.17 provide a simple recurrence for the singular cohomology of uniform matroids. In this subsection, we use this recurrence to produce closed formulas for the refined Hodge–Poincaré polynomials.

Recall the truncation exact sequence

$$\cdots \rightarrow \bigoplus_{H \in \mathcal{H}(U_{r,n})} H_i(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_j \rightarrow H_i(K(\mathbb{Q}[\Sigma_{U_{r,n}}]))_{j+1} \rightarrow H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))_{j+1} \rightarrow \cdots$$

with connecting homomorphism

$$\delta_i: H_i(K(\mathbb{Q}[\Sigma_{U_{r-1,n}}]))_j \rightarrow \bigoplus_H H_{i-1}(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_{j-1}.$$

Lemma 3.17 shows that δ_i is injective for $i \geq 1$, while δ_0 vanishes trivially.

In order to get a recurrence for the refined Hodge–Poincaré polynomials, we first express the dimensions of $H_i(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_j$ in terms of *Eulerian polynomials*.

Definition 4.16. The n th Eulerian polynomial $A_n(x)$ is the generating function for permutations in \mathfrak{S}_n with respect to the number of excedences:

$$A_n(x) = \sum_{\sigma \in \mathfrak{S}_n} x^{\text{exc}(\sigma)},$$

and we set $A_0(x) = 1$ by convention.

Lemma 4.17. For $i \geq 0$,

$$\sum_{H \in \mathcal{H}(U_{r,n})} \sum_{j=0}^{r-2} \dim H_i(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_j \cdot x^j = \binom{n}{r-1} \binom{n-r}{i} A_{r-1}(x).$$

Proof. Lemma 3.14 implies that

$$\sum_{j=0}^{r-2} \dim H_i(K(\mathbb{Q}[\overline{\text{St}}(\rho_H)]))_j \cdot x^j = \binom{n-r}{i} \underline{H}_{U_{r-1,r-1}}(x),$$

for each $H \in \mathcal{H}(U_{r,n})$. It is well-known that $\underline{H}_{U_{r-1,r-1}}(x) = A_{r-1}(x)$ (see [HRS21, Theorem 5.1]). As $|\mathcal{H}(U_{r,n})| = \binom{n}{r-1}$, the lemma follows. \square

For $i \geq 1$, the truncation exact sequence therefore provides the recurrence

$$\underline{H}_{U_{r,n}}^i(x) = \binom{n}{r-1} \binom{n-r}{i} A_{r-1}(x) \cdot x - \underline{H}_{U_{r-1,n}}^{i+1}(x) \cdot x,$$

which produces the following formula.

Proposition 4.18. For $i \geq 1$,

$$\underline{H}_{U_{r,n}}^i(x) = (-1)^{r-1} \binom{n-1}{i+r-1} \cdot x^{r-1} + \sum_{j=1}^{r-1} (-1)^{r+1-j} \binom{n}{j} \binom{n-j-1}{i+r-j-1} A_j(x) \cdot x^{r-j}.$$

Proof. The base case is $\underline{H}_{U_{1,n}}^i(x) = \binom{n-1}{i}$, and the result follows from induction on r . \square

The case $i = 0$ is for the Chow polynomial, and we have the recurrence

$$\underline{H}_{U_{r,n}}(x) = \binom{n}{r-1} A_{r-1}(x) \cdot x + \underline{H}_{U_{r-1,n}}(x) - \underline{H}_{U_{r-1,n}}^1(x) \cdot x,$$

which recovers the following formula of Ferroni et al.

Proposition 4.19 ([Fer+24, Equation (13)]). For $r \geq 1$,

$$\underline{H}_{U_{r,n}}(x) = \sum_{j=0}^{r-1} \sum_{m=0}^{r-j-1} (-1)^m \binom{n}{j} \binom{n-j-1}{m} A_j(x) \cdot x^{r-m-j-1}.$$

Proof. By induction and Proposition 4.18, one can show that

$$\underline{H}_{U_{r,n}}(x) = \sum_{m=0}^{r-1} (-1)^m \binom{n-1}{m} \cdot x^m + \sum_{j=1}^{r-1} \sum_{m=0}^{r-j-1} (-1)^m \binom{n}{j} \binom{n-j-1}{m} A_j(x) \cdot x^{m+1}.$$

The desired equation follows by taking the reverse polynomial of each side and using the fact $\underline{H}_{U_{k,n}}(x)$ and $A_j(x)$ are palindromic. \square

Example 4.20. We use these formulas to list the Betti diagrams for the singular cohomology of loopless uniform matroids on ground set [5] in Table 1.

$U_{1,5}$	0	1	2	3	4		$U_{2,5}$	0	1	2	3		$U_{3,5}$	0	1	2
0	1	4	6	4	1		0	1	0	0	0		0	1	0	0
							1	1	9	11	4		1	11	20	10
													2	1	9	6

$U_{4,5}$	0	1					$U_{5,5}$	0
0	1	0					0	1
1	21	10					1	26
2	21	30					2	66
3	1	4					3	26
							4	1

TABLE 1. Betti diagrams for the singular cohomology of loopless uniform matroids on ground set [5].

5. LEFSCHETZ PROPERTIES

In this final section, we show that the singular cohomology ring of a uniform matroid satisfies a slight weakening of Hard Lefschetz which we call the *quasi-projective strong Lefschetz property*.

5.1. Quasi-projective Lefschetz properties. We begin by recalling the weak and strong Lefschetz properties for standard graded Artinian algebras. See [MN13] for a survey.

Definition 5.1. Let A^\bullet be an Artinian graded algebra and $\ell \in A^1$ a generic linear form.

- (1) We say that A^\bullet satisfies the *weak Lefschetz property* (WLP) if

$$\times \ell: A^i \longrightarrow A^{i+1}$$

has maximal rank for all $i \in \mathbb{Z}_{\geq 0}$.

- (2) We say that A^\bullet satisfies the *strong Lefschetz property* (SLP) if

$$\times \ell^d: A^i \longrightarrow A^{i+d}$$

has maximal rank for all $i, d \in \mathbb{Z}_{\geq 0}$.

For a smooth, quasi-projective variety X , the singular cohomology ring $H^\bullet(X)$ is Artinian, so one can ask if it satisfies WLP or SLP. However, this will often fail when torus factors are involved.

Example 5.2. Let Y be a smooth, projective toric variety of (complex) dimension s , and consider $X = Y \times (\mathbb{C}^*)^t$.

Hodge theory implies that for a generic $\ell \in \text{Gr}_2^W H^2(Y)$, the map

$$\times \ell^d: H^k(Y) \longrightarrow H^{k+2d}(Y)$$

is injective for $d \leq s - k$ and surjective for $d \geq s - k$. Therefore, $H^\bullet(Y)$ satisfies SLP.

However, $H^\bullet(X)$ does not satisfy SLP when $t \geq 2$. Indeed, $H^k(X) \cong \bigoplus_{i=0}^t H^{k-i}(Y)^{\oplus \binom{t}{i}}$, so $\times \ell^{s-k+1}: H^k(X) \rightarrow H^{2s-k+2}(X)$ is neither injective on the subfactor $H^k(Y) \rightarrow H^{2s-k+2}(Y)$ nor surjective on the subfactor $H^{k-2}(Y)^{\oplus \binom{t}{2}} \rightarrow H^{2s-k}(Y)^{\oplus \binom{t}{2}}$ for $2 \leq k \leq s$.

The weight filtration rectifies the situation. Now, $\text{Gr}_j^W H^k(X) \cong H^{2k-j}(Y)^{\oplus \binom{t}{j-k}}$, so

$$\times \ell^d: \text{Gr}_j^W H^k(X) \rightarrow \text{Gr}_{j+2d}^W H^{k+2d}(X)$$

is injective for $d \leq s + j - 2k$ and surjective for $d \geq s + j - 2k$.

This example motivates using the weight filtration to define Lefschetz properties for quasi-projective varieties.

Definition 5.3. Let X be a smooth, quasi-projective variety and $\ell \in \text{Gr}_2^W H^2(X)$ a generic hyperplane class in the singular cohomology ring.

(1) We say that $H^\bullet(X)$ satisfies the *quasi-projective weak Lefschetz property* (qpWLP) if

$$\times \ell: \text{Gr}_j^W H^k(X) \longrightarrow \text{Gr}_{j+2}^W H^{k+2}(X)$$

has full rank for all $j, k \in \mathbb{Z}_{\geq 0}$.

(2) We say that $H^\bullet(X)$ satisfies the *quasi-projective strong Lefschetz property* (qpSLP) if

$$\times \ell^d: \text{Gr}_j^W H^k(X) \longrightarrow \text{Gr}_{j+2d}^W H^{k+2d}(X)$$

has full rank for all $d, j, k \in \mathbb{Z}_{\geq 0}$.

5.2. Lefschetz in the singular cohomology ring of a uniform matroid. We now show that the singular cohomology ring of a uniform matroid satisfies the quasi-projective strong Lefschetz property.

Our proof uses the truncation exact sequence (2) to bootstrap Lefschetz properties for the singular cohomology of $U_{r,n}$ from the singular cohomology of the permutohedral variety.

Lemma 5.4. *For a hyperplane $H \in \mathcal{L}(U_{r,n})$ and generic $\ell \in \text{Gr}_2^W H^2(X_{\overline{\text{St}}(\rho_H)})$, the map*

$$\times \ell^d: \text{Gr}_j^W H^k(X_{\overline{\text{St}}(\rho_H)}) \rightarrow \text{Gr}_{j+2d}^W H^{k+2d}(X_{\overline{\text{St}}(\rho_H)})$$

is injective for $0 \leq d \leq r + j - 2k - 2$ and surjective for $d \geq r + j - 2k - 2$.

Proof. By [Bin26, Lemma A.3], $H^\bullet(X_{\overline{\text{St}}(\rho_H)}) \cong H^\bullet(X_{\text{St}(\rho_H)})$, so it suffices to consider $X_{\text{St}(\rho_H)}$. Now, $X_{\text{St}(\rho_H)} \cong X_{\Sigma_{U_{r-1,r-1}}} \times (\mathbb{C}^*)^{n-|H|-1}$. As $X_{\Sigma_{U_{r-1,r-1}}}$ is the permutohedral variety of dimension $r - 2$, the result follows as in Example 5.2. \square

Now consider an arbitrary class $\ell \in \text{Gr}_2^W H^2(X_{\Sigma_{U_{r,n}}})$, and let ℓ_H denote its pull-back to $X_{\overline{\text{St}}(\rho_H)}$ and $\bar{\ell}$ its pull-back to $X_{\Sigma_{U_{r-1,n}}}$. In terms of linear forms in the Stanley–Reisner ring, ℓ_H quotients out the indeterminates x_F such that $F \not\leq H$ and $\bar{\ell}$ quotients out the indeterminates x_H for $H \in \mathcal{H}(U_{r,n})$. The truncation exact sequence (2) produces commuting squares

$$\begin{array}{ccc} \text{Gr}_j^W H^k(\Sigma_{U_{r-1,n}}) & \xrightarrow{\delta} & \bigoplus_{H \in \mathcal{H}(U_{r,n})} \text{Gr}_{j-2}^W H^{k-1}(X_{\overline{\text{St}}(\rho_H)}) \\ \downarrow \times \bar{\ell}^d & & \downarrow \oplus \times \ell_H^d \\ \text{Gr}_{j+2d}^W H^{k+2d}(\Sigma_{U_{r-1,n}}) & \xrightarrow{\delta} & \bigoplus_{H \in \mathcal{H}(U_{r,n})} \text{Gr}_{j+2d-2}^W H^{k+2d-1}(X_{\overline{\text{St}}(\rho_H)}) \end{array}$$

and

$$\begin{array}{ccc} \bigoplus_{H \in \mathcal{H}(U_{r,n})} \text{Gr}_{j-2}^W H^{k-2}(X_{\overline{\text{St}}(\rho_H)}) & \longrightarrow & \text{Gr}_j^W H^k(X_{\Sigma_{U_{r,n}}}) \\ \downarrow \oplus \times \ell_H^d & & \downarrow \times \ell^d \\ \bigoplus_{H \in \mathcal{H}(U_{r,n})} \text{Gr}_{j+2d-2}^W H^{k+2d-2}(X_{\overline{\text{St}}(\rho_H)}) & \longrightarrow & \text{Gr}_{j+2d}^W H^{k+2d}(X_{\Sigma_{U_{r,n}}}). \end{array}$$

Theorem 5.5. *Let $\ell \in \text{Gr}_2^W H^2(X_{\Sigma_{U_{r,n}}})$ be generic. Then the map*

$$\times \ell^d : \text{Gr}_j^W H^k(X_{\Sigma_{U_{r,n}}}) \rightarrow \text{Gr}_{j+2d}^W H^{k+2d}(X_{\Sigma_{U_{r,n}}})$$

*is injective for $0 \leq d \leq r + j - 2k - 1$ and surjective for $d > r + j - 2k - 1$. In particular, the singular cohomology ring of a uniform matroid satisfies *qpSLP*.*

Proof. If $j = k$ or $r = n$, this is a weaker statement than Hard Lefschetz for the Chow ring of a matroid or the permutohedral variety, respectively. Therefore, assume $j \neq k$ and $r < n$.

When $0 \leq d \leq r + j - 2k - 1$, we use the first commuting square from above. We may find $\hat{\ell} \in \text{Gr}_2^W H^k(X_{\Sigma_{U_{r+1,n}}})$ such that $\ell = \hat{\ell}$. By genericity, we may also assume each $\hat{\ell}_H$ is generic. The map $\oplus \times \hat{\ell}_H^d$ is injective by Lemma 5.4. The horizontal maps are also injective by Lemma 3.17, so $\times \ell^d$ is injective as well.

When $d > r + j - 2k - 1$, we use the second square. The map $\oplus \times \ell_H^d$ is surjective by Lemma 5.4, and the horizontal maps are surjective by Lemma 3.17. This implies $\times \ell^d$ is surjective. \square

Remark 5.6. This proof can be adapted to show that the conclusion of Theorem 5.5 holds for any $\ell \in \text{Gr}_2^W H^2(X_{\Sigma_{U_{r,n}}})$ which is the pull-back of a class $\hat{\ell}$ from the *ample cone* of $\Sigma_{U_{r+1,n}}$ (see [AHK18, Definition 4.2]). The idea is that both $\hat{\ell}_H$ and $\tilde{\ell}_H$ are in the ample cone of $\text{St}(\rho_H)$ for each $H \in \mathcal{H}(U_{r+1,n})$ and $H \in \mathcal{H}(U_{r,n})$, respectively, by [AHK18, Proposition 4.4]. Then [AHK18, Theorem 8.8] implies the conclusion of Lemma 5.4 holds for $\hat{\ell}_H$ and $\tilde{\ell}_H$, and the rest of the proof follows verbatim.

A combinatorial consequence of the quasi-projective strong Lefschetz property is the following.

Corollary 5.7. *The refined Hodge–Poincaré polynomials $\underline{H}_{U_{r,n}}^i(x)$ have unimodal coefficients. Equivalently, the Betti diagram for the Hodge numbers of $U_{r,n}$ has unimodal columns.*

In fact, we conjecture that a stronger statement is true.

Conjecture 5.8. *The refined Hodge–Poincaré polynomials $\underline{H}_{U_{r,n}}^i(x)$ are real-rooted.*

Real-rootedness is known for the Chow polynomials by [BV25, Theorem 1.1]. As evidence for the other refined Hodge–Poincaré polynomials, we have verified each $\underline{H}_{U_{r,n}}^i(x)$ has ultra-log-concave coefficients for $n \leq 230$ using Proposition 4.18 and a short Python script.

REFERENCES

- [AHK18] Karim Adiprasito, June Huh, and Eric Katz. “Hodge theory for combinatorial geometries”. In: *Ann. of Math. (2)* 188.2 (2018), pp. 381–452.
- [Bin26] Kyle Binder. *The Singular Cohomology Ring of a Matroid*. 2026. arXiv: 2412.05732 [math.CO].
- [BV25] Petter Brändén and Lorenzo Vecchi. *Chow polynomials of uniform matroids are real-rooted*. 2025. arXiv: 2501.07364 [math.CO].
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*. Vol. 124. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2011, pp. xxiv+841.
- [Fer+24] Luis Ferroni et al. “Hilbert–Poincaré series of matroid Chow rings and intersection cohomology”. In: *Adv. Math.* 449 (2024), Paper No. 109733, 55.
- [FF24] Matthias Franz and Xin Fu. “Cohomology of smooth toric varieties: naturality”. In: *J. Pure Appl. Algebra* 228.6 (2024), Paper No. 107590, 38.

- [FY04] Eva Maria Feichtner and Sergey Yuzvinsky. “Chow rings of toric varieties defined by atomic lattices”. In: *Invent. Math.* 155.3 (2004), pp. 515–536.
- [Hos26] Elena Hoster. “The Chow and Augmented Chow Polynomials of Uniform Matroids”. In: *Electron. J. Combin.* 33.1 (2026), P1.36.
- [HRS21] Thomas Hameister, Sujit Rao, and Connor Simpson. “Chow rings of vector space matroids”. In: *J. Comb.* 12.1 (2021), pp. 55–83.
- [JMS19] Martina Juhnke-Kubitzke, Satoshi Murai, and Richard Sieg. “Local h -vectors of quasi-geometric and barycentric subdivisions”. In: *Discrete Comput. Geom.* 61.2 (2019), pp. 364–379.
- [MN13] Juan Migliore and Uwe Nagel. “Survey article: a tour of the weak and strong Lefschetz properties”. In: *J. Commut. Algebra* 5.3 (2013), pp. 329–358.
- [Oxl11] James Oxley. *Matroid theory*. Second. Vol. 21. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, 2011, pp. xiv+684.
- [Tot14] Burt Totaro. “Chow groups, Chow cohomology, and linear varieties”. In: *Forum Math. Sigma* 2 (2014), Paper No. e17, 25.
- [Web04] Andrzej Weber. “Weights in the cohomology of toric varieties”. In: *Cent. Eur. J. Math.* 2.3 (2004), pp. 478–492.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY
Email address: kbinde1@lsu.edu